The Reduction of Waste in the Operating Room via an Intraoperative Recycling Initiative



Catherine A. Ruggiero, SRNA, BSN, BS, RN¹; Dayna M. Dombrowsky, SRNA, BSN, RN²; Rachel A. Bosco, SRNA, BSN, RN³

University of Pennsylvania School of Nursing, Philadelphia, PA



Background & Purpose

- United States (US) healthcare facilities produce 6,600 tons of waste daily (Lee et al., 2012)
- From 2010-2018, the US healthcare industry contributed to a 6% increase in total greenhouse gas emissions (Eckelman et al., 2020).
- ~40-70% of healthcare waste is generated within the OR (Yeoh et al., 2020)
- Anesthesia care contributes to 25% of intraoperative material wastes (Yeoh et al., 2020)
- Purpose: Educate the intraoperative team on environmentally sustainable practices and to increase the amount of recycled waste.

Methods

- Project Design: Quality Improvement (QI)
- Setting: Large urban community teaching hospital
- *Participants*: Anesthesiologists, CRNAs, Anesthesia techs, SRNAs, OR RNs, Surgical techs, and EVS
- Measures: 1.) Staff knowledge and 2.) waste volume vs recycled volume in pounds.
- 2-week pre-intervention baseline trash assessment
- Knowledge Assessment Tool (Pre-Education)
- Educational PowerPoint presentation

SCAN MF

- Knowledge Assessment Tool (Post-Education)
- 4-week pilot recycling program with education reinforcements such as a QR code (see below) that links to our website, Team Recycle.

Please don't forget

to recycle!

Scan the OR code to get


more information on

what you can recycle in

this OR

Results

- Measure 1 Results: The mean number of correctly answered questions pre-education was 60.3% and post-education was 72.6% (p-value <0.0181).
- Measure 2 Results: A total of 1,369.09lbs of waste and 253.22lbs of recyclable material was collected during 4-weeks.
- Mean amount of unsorted waste = 23.61lbs/day
- Mean amount of recyclable material = 4.29lbs/day





Summary & Conclusion

- 15.6% decrease in volume of waste placed in the unsorted trash.
- 100% increase in recycling weight after the completion of the 4-week pilot program.
- It cannot be assumed with complete certainty that the reduction of waste generation directly correlates to the intraoperative recycling pilot program.
- The increase in the amount of material placed in the recycling receptacles demonstrates that our educational intervention was successfully translated into practice and was sustained over a 4-week period.

Acknowledgements

 We would like to thank our faculty lead, Dr. Susan Renz, PhD, DNP, GNP-BC, and our site leads Tracy Flood, MSN, CRNA, and Dr. Angelarosa DiDonato, DNP, CRNA

